

**Huntington 2018 - Tick Infection Rates**  
**Collection site is located within the 11743 zip code**

Infection Rate (%)	Pathogen	Disease	Stage	Tick
62	<i>B. burgdorferi</i>	Lyme Disease	Adult	Blacklegged Tick
28	<i>B. microti</i>	Babesiosis	Adult	Blacklegged Tick
23.5	<i>B. burgdorferi</i>	Lyme Disease	Nymph	Blacklegged Tick
17.6	<i>B. microti</i>	Babesiosis	Nymph	Blacklegged Tick
16	<i>A. phagocytophilum</i>	Anaplasmosis	Adult	Blacklegged Tick
6	<i>B. miyamotoi</i>	Relapsing Fever	Adult	Blacklegged Tick
2	<i>A. phagocytophilum</i>	Anaplasmosis	Nymph	Blacklegged Tick
2	<i>B. miyamotoi</i>	Relapsing Fever	Nymph	Blacklegged Tick
2	Powassan virus	Powassan Virus Disease	Adult	Blacklegged Tick
1.9	Powassan virus	Powassan Virus Disease	Nymph	Blacklegged Tick

\*While nymphal ticks often have lower infection rates than adult ticks, they are typically responsible for causing more disease in humans, as their smaller size allows them to avoid notice